2018

CHEMISTRY

(Major)

Paper: 6.2

(Physical Chemistry)

Full Marks: 60

Time: 3 hours

The figures in the margin indicate full marks for the questions

1. Answer the following in brief:

 $1 \times 7 = 7$

- (a) The ionic radii of Cs⁺ and Cl⁻ ions are 1.69 Å and 1.81 Å respectively. Predict the coordination number of Cs⁺.
- (b) Yttrium barium copper oxide superconductor is often referred to as the 123 superconductor. Why?
- (c) Arrange the following in increasing order of their effectiveness in coagulating ferric hydroxide sol:

 Na₂SO₄, KCl, K₃[Fe(CN)₆]

- (d) When a freshly prepared precipitate of Fe(OH)₃ is treated with water and a small amount of FeCl₃ solution, Fe(OH)₃ is converted to colloidal solution. What is the role of FeCl₃ in this process?
- (e) Why should one always use purest monomer in free-radical polymerization?
- (f) What do you mean by dominant configuration?
- (g) The weight average and number average molecular weight of a polymer is 60000 kg mol⁻¹ and 40000 kg mol⁻¹ respectively. What will be the polydispersity index of the polymer?

2. Answer the following:

 $2 \times 4 = 8$

(a) Lithium borohydride, LiBH₄ crystallizes in an orthorhombic system with 4 molecules per unit cell. The unit cell dimensions are a = 6.81 Å, b = 4.43 Å and c = 7.17 Å. If its molar mass is 21.76 g mol⁻¹, calculate the density of the crystal.

- (b) Give the physical significance of molecular partition function.
- (c) Explain the difference between accuracy and precision with examples.
- (d) 100 ml of a colloidal solution is completely precipitated by addition of 5 ml of 1 M NaCl solution. Calculate the coagulation value of NaCl.
- 3. (a) How does electrical conductivity of a metal and a semiconductor vary with temperature? What do you mean by n-type and p-type semiconductors? Explain with appropriate diagram and example.

 1+4=5

Or

What are Schottky and Frenkel defects?

Derive an expression for the number of
Schottky defects in a crystal. 2+3=5

(b) Deduce an expression for the entropy of monoatomic perfect gas in terms of partition function.

5

Derive an expression for rotational partition function. rotational The constant of gaseous HCl, determined from microwave spectroscopy is 10.59 cm⁻¹. Calculate the rotational partition function of HCl at 100 K. 3+2=5

Distinguish between repeatable and reproducible results. Analyzing of a sample of iron ore gave the following percentage values for the iron content:

> 7.08, 7.21, 7.12, 7.09, 7.16, 7.14, 7.07, 7.14, 7.18, 7.11

Calculate the mean, standard deviation and coefficient of variations for the values. 1+4=5

- 4. Answer either (a), (b) and (c) or (d), (e) and (f):
 - A reflection from the (111) planes of a cubic crystal was observed at a glancing angle of 11·2° when CuK_α X-rays of wavelength 154 pm were used. What is the length of the side of the unit cell?
 - Show that for an atom to occupy a tatrahedral void, its radius must be 0.225 times the radius of the sphere.

Non-stoichiometric cuprous oxide, Cu2O can be prepared in laboratory. In this oxide, copper to oxygen ratio is slightly less than 2:1. How will you account for the fact that this substance is a p-type semiconductor on the basis of the above stated information?

3

2

- (d) What is radius-ratio? How does radiusratio help in determining the structure of ionic solids and coordination number of ions? Explain. 1+3=4
- Why does zinc oxide exhibit enhanced electrical conductivity on heating?
- Explain the following terms with examples: $2 \times 2 = 4$
 - (i) Ferrimagnetism
 - (ii) Piezoelectricity
- 5. Answer either (a), (b) and (c) or (d), (e) and (f):
 - (a) Discuss the osmotic pressure method for determination of molar mass of polymers. Why does this method give number average molar mass only? 4+1=5
 - The intrinsic viscosity of myosin is 217 cm³ g⁻¹. Calculate the appropriate concentration of myosin in water if it has a relative viscosity of 1.5.

8A/894

(Continued)

3

8A/894

(Turn Over)

3

- (c) What are lyophilic and lyophobic sols? Give one example of each. 1+1=2
- (d) Discuss the kinetics of condensation polymerization. Give an example of a polymer produced by this method. 3+1=4
- In a polymer sample, 30% molecules have a molar mass 20000, 40% have molar mass 60000 and the rest have 30000. Calculate weight average and number average molar mass of the polymer.
- Account for the origin of charge on 3 colloidal particles in detail.
- 6. Answer either (a) and (b) or (c) and (d):
 - (a) Derive the Boltzmann distribution law. Give its physical significance. 5+1=6
 - (b) Distribute three energy quanta among three particles. Calculate the probability of each distribution.
 - (c) Using the concept of partition function, deduce an expression for the internal energy of a monoatomic perfect gas.

Hence find an expression for the heat capacity at constant volume. Graphically show how heat capacity of diatomic molecules varies with temperature. 3+2+1=6

For a diatomic molecule vibrating as a simple harmonic oscillator, obtain an expression for vibrational partition function.

8A-4500/894

3